

Study questions

1. Define and use the Rankine-Hugoniot jump condition to compute the shock speed for the following problem

$$\begin{aligned}
 u_t + uu_x &= 0 & -\infty < x < \infty, \quad t > 0 \\
 u(x, 0) &= \begin{cases} 1 & x \leq 0 \\ 0 & \text{otherwise} \end{cases}
 \end{aligned}$$

2. Define the entropy condition for a scalar conservation law.

$$u_t + f(u)_x = 0 \quad -\infty < x < \infty, \quad t > 0$$

with a convex flux function $f(u)$. The shock is moving with speed s and the state to the left is given by u_L and the state to the right by u_R .

Why do we need an entropy condition ?

3. Define a total variation decreasing (TVD) method. Why is this a desirable property ?
4. Investigate the one-sided difference scheme

$$u_j^{n+1} = u_j^n - a \frac{\Delta t}{\Delta x} (u_j^n - u_{j-1}^n)$$

for the advection equation

$$u_t + au_x = 0$$

Consider the cases $a > 0$ and $a < 0$.

- a) Prove that the scheme is consistent and find the order of accuracy. Assume k/h constant.
 - b) Determine the stability requirement for $a > 0$ and show that it is unstable for $a < 0$.
5. Apply Lax-Friedrichs scheme to the linear wave equation

$$u_t + au_x = 0.$$

- a) Write down the modified equation.
- b) What type of equations is this?
- c) What kind of behavior can we expect from the solution?

6. A the three-point centered scheme applied to

$$u_t + au_x = 0, \quad a > 0.$$

yields the approximation

$$u_j^{n+1} = u_j^n + \frac{a\Delta t}{2\Delta x}(u_{j+1} - u_{j-1})$$

Show that this approximation is not stable even though the CFL condition is fulfilled.

7. What does Lax's equivalence theorem state?
 8. What is the condition on the $n \times n$ real matrix $A(\mathbf{u})$ for the system

$$\mathbf{u}_t + A\mathbf{u}_x = 0$$

to be hyperbolic ?

9. The barotropic gas dynamic equations

$$\begin{aligned} \rho_t + \rho u_x &= 0 \\ u_t + uu_x + \frac{1}{\rho}p_x &= 0 \end{aligned} \tag{1}$$

where

$$p = p(\rho) = C\rho^\gamma$$

i and C a constant, can be linearized by considering small perturbations (ρ', u') around a motionless gas.

- a) Let $\rho = \rho_0 + \rho'$ and $u = u_0 + u'$ where $u_0 = 0$. Linearize the system (1) and show that this yields the following linear system (the primes has been dropped)


$$\begin{aligned} \rho_t + \rho_0 u_x &= 0 \\ u_t + \frac{a^2}{\rho_0} \rho_x &= 0 \end{aligned} \tag{2}$$

where a is the speed of sound. a and ρ_0 are constants.

- b) Is the system given by (2) a hyperbolic system? Motivate your answer.
 c) Determine the characteristic variables in terms of ρ and u .
 d) Determine the partial differential equations the characteristic variables fulfill - characteristic formulation.
 e) Given initial conditions at $t = 0$ and let $-\infty < x < \infty$ (no boundaries)

$$\rho(0, x) = \sin(x) \quad u(0, x) = 0$$

determine the analytical solution to (2) for $t > 0$. Hint: Start from the characteristic formulation.



10. The linearized form of (2) is given by

$$\begin{pmatrix} \rho \\ u \end{pmatrix}_t + \underbrace{\begin{pmatrix} 0 & \rho_0 \\ a^2/\rho_0 & 0 \end{pmatrix}}_A \begin{pmatrix} \rho \\ u \end{pmatrix}_x = 0, \quad (3)$$

where a is the speed of sound. a and ρ_0 are constants.

a) Draw the domain of dependence of the solution to the system (3) in a point P in the x-t plane.

b) The system is solved numerically on a grid given by $x_j = j\Delta x, j = 0, 1, 2, \dots$ and $t_n = n\Delta t, n = 0, 1, 2, \dots$ using an explicit three-point scheme, see the figure below.

Draw the domain of dependence of the numerical solution at P (in the same figure as a)) of the three-point scheme in the case when

- i) the CFL condition is fulfilled
- ii) the CFL condition is NOT fulfilled.

Assume that P is a grid point.

11. To solve Euler equations in 1D

$$\begin{aligned} \rho_t + \rho u_x + u \rho_x &= 0 \\ u_t + uu_x + \frac{1}{\rho} p_x &= 0 \\ p_t + \rho c^2 u_x + u p_x &= 0 \end{aligned}$$

How many boundary conditions must be added at (motivate your answer) inflow boundary when the flow is

- a) Supersonic
- b) Subsonic

outflow boundary when the flow is

- c) Supersonic
- d) Subsonic

12. a) Show that a vector field w_i can be decomposed into

$$w_i = u_i + \frac{\partial p}{\partial x_i}$$

where u is divergence free and parallel to the boundary.

b) Apply this to the Navier-Stokes equations, show that the pressure term disappears and recover an equation for the pressure from the gradient part.

13. From the differential form of the Navier-Stokes equations obtain
 - a) the Navier-Stokes equations in integral form used in finite-volume discretizations,
 - b) a variational form of the Navier-Stokes equations used in finite-element discretizations.
14. a) Write down the appropriate function spaces for the pressure and velocity used to define weak solutions to the Navier-Stokes equations.
 - b) Explain the concept of essential and natural boundary conditions.
15. a) How is a finite-element approximation defined?
 - b) Explain how to convert a FEM discretization to an algebraic problem, e.g. that the Navier-Stokes equations yield

$$\begin{pmatrix} \nu \mathbf{K} + \mathbf{C}(u) & \mathbf{G} \\ \mathbf{G}^T & \mathbf{0} \end{pmatrix} \begin{pmatrix} u \\ p \end{pmatrix} = \begin{pmatrix} f \\ 0 \end{pmatrix}$$

16. a) By choosing $\mathbf{z}_h = \mathbf{u}_h$ in the finite element approximation of the Stokes problem, show that any Galerkin velocity solution is stable.
 - b) Describe and interpret the LBB condition.
17. Choice of elements. Discrete elements pairs
 - a) constant pressure-bilinear velocities,
 - b) the Taylor-Hood pair,
 - c) a stable choice with piecewise linear velocities.
18. a) Derive the finite volume (FV) discretization on arbitrary grids of the continuity equation ($\partial u_i / \partial x_i = 0$),
 - b) derive the FV discretization for Laplace equation on a Cartesian grid,
 - c) show that both are equivalent to a central difference approximation for Cartesian grids.
19. Derive the finite element (FEM) discretization for Laplace equation on a Cartesian grid and show that it is equivalent to a central difference approximation.

20. Iterative techniques for linear systems.
- Define Gauss-Seidel iterations for the Laplace equations,
 - Define the 2-level multigrid method for the Laplace equation,
21. State the difficulties associated with the the finite-volume discretizations of the Navier-Stokes equations on a colocated grid? and show the form of the spurious solution which exist.
22. a) Define an appropriate staggered grid that can be used for the discretization of the Navier-Stokes equations,
- write down the FV discretization of the Navier-Stokes equations on a staggered cartesian grid,
 - discuss how to treat noslip and inflow/outflow boundary conditions.
23. Time dependent flows.
- Define a simple projection method for the time dependent Navier-Stokes equations

$$\frac{d}{dt} \begin{pmatrix} u \\ 0 \end{pmatrix} + \begin{pmatrix} \mathbf{N}(u) & \mathbf{G} \\ \mathbf{D} & \mathbf{0} \end{pmatrix} \begin{pmatrix} u \\ p \end{pmatrix} = \begin{pmatrix} f \\ g \end{pmatrix}$$

- show in detail the equation for the pressure to be solved at each time step and discuss the boundary conditions for the pressure.
24. Time step restriction for Navier-Stokes solutions.
- Motivate the use of an appropriate form of the advection-diffusion equation as a model equation for stability analysis,
 - derive the time step restrictions for the 1D version of that equation,
 - state the 2D equivalent of that restriction.